Current Trends In The Biology Of Orthodontic Tooth Movement: Cellular, Molecular, And Translational Perspectives

Authors

K. Riteesh Dani[1], Y. Muni Akash[1], Harshitha Nagaram[2*], S.V. Kala Vani[3] , Hemadri.S[4]

1.Undergraduate, 2. Senior Lecturer, 3. Head of Department, 4. Reader

Department Of Orthodontics And Dentofacial Orthopaedics CKS Teja Institute Of Dental Sciences And Research, Tirupati-517501, Andhra Pradesh, India.

Corresponding Author*

Dr.Harshitha Nagaram

Senior Lecturer

Department Of Orthodontics And Dentofacial Orthopaedics, CKS Teja Institute Of Dental Sciences And Research, Tirupati-517501, Andhra Pradesh, India.

Received: Oct 01, 2026 Accepted: Oct 28, 2026 Published: Nov 03, 2026

ABSTRACT

Orthodontic tooth movement (OTM) is a biologically mediated process initiated by mechanical forces on teeth, resulting in remodelling of the periodontal ligament (PDL) and alveolar bone. Recent advances in cellular and molecular biology have substantially deepened understanding of the mechanobiological mechanisms underlying OTM. Current research emphasises osteocytes' central role as mechanosensors, the importance of inflammatory and immune responses, and the elaborate signalling networks that regulate bone resorption and formation. Notable emerging areas include autophagy, extracellular matrix remodelling, chronobiology, systemic immune interactions, and translational approaches to accelerate tooth movement. This review integrates contemporary evidence on the biological basis of OTM, focusing on recent developments and future directions that may improve the efficiency, predictability, and biological safety of orthodontic treatment.

Keywords: Orthodontic tooth movement; mechanobiology; periodontal ligament; osteocytes; RANKL; bone remodeling

INTRODUCTION

Orthodontic tooth movement is the result of the biological response of the periodontal and alveolar tissues to the application of mechanical forces. It is not simply a matter of mechanical forces, it is the result of mechanotransduction, inflammatory signalling and coordinated bone remodelling.[1-3] The classical pressure–tension theory describes bone resorption on the compression side and bone formation on the tension side during OTM. Recent molecular biology research reveals that complex cellular interactions among periodontal ligament fibroblasts, osteoblasts, osteoclasts, osteocytes, endothelial cells, and immune cells mediate this process.[4-6] A greater understanding of these functions is necessary to improve treatment efficiency, reduce adverse effects such as root resorption, and advance biologically based orthodontic therapies.[7,8]

BIOLOGICAL BASIS OF ORTHODONTIC TOOTH MOVEMENT

The PDL is the main site of force sensing during orthodontic treatment. Mechanical loading has an effect on the vascularity, ECM (extracellular matrix) organisation and cellular activity of the PDL.[9-10]In response to mechanical strain the PDL fibroblasts will secrete cytokines, prostaglandins and growth factors which in turn initiate the bone remodelling process.[11] Too much force can lead to local ischaemia and hyalinisation and thus impede tooth movement, whereas the ideal amount of force will result in continuous, efficient remodelling.[12]

Alveolar bone remodelling is the result of tightly coupled osteoclastic resorption and osteoblastic formation. Osteoclasts resorb bone on the compression side and osteoblasts form new bone on the tension side, thus maintaining the structural integrity of the periodontium.[13,14]

MECHANOTRANSDUCTION AND CELLULAR MECHANISMS

Mechanotransduction is the process of converting mechanical force into a biochemical signal. PDL fibroblasts and osteoblasts are all capable of mechanotransduction but osteocytes are now considered to be bone's principal mechanosensors.[15-17]Osteocytes regulate bone remodelling by modulating the production of a range of signalling molecules (RANKL, OPG, sclerostin, nitric oxide and prostaglandins) in response to mechanical strain.[18-20] These molecules in turn influence osteoclast and osteoblast activity and so osteocytes are key regulators of the response to orthodontic force.[21]

MOLECULAR SIGNALLING PATHWAYS IN TOOTH MOVEMENT

The RANK/RANKL/OPG axis is of fundamental importance in osteoclastogenesis during OTM. RANKL (which is produced by PDL cells, osteoblasts and osteocytes) binds to RANK on the osteoclast precursor and stimulates bone resorption.[22,23] OPG functions as a decoy receptor and thus inhibits the RANKL/RANK interaction. The RANKL/OPG ratio is therefore a key determinant of the rate of tooth movement.[24]

Orthodontic force leads to a sterile inflammatory response (increased IL-1β, IL-6, TNF-α and PGE2).[25-27]This response promotes osteoclast recruitment, increases vascular permeability and accelerates extracellular matrix turnover.

The Wnt/beta-catenin pathway is the principal regulator of osteoblast differentiation and bone formation. Mechanical loading leads to a reduction in osteocyte-derived sclerostin which in turn activates Wnt signalling.[28,29] Other pathways (MAPK, NF-kappaB and TGF-beta signalling) also play a role in the cellular adaptation that occurs during OTM.[30]

EMERGING TRENDS IN THE BIOLOGY OF ORTHODONTIC TOOTH MOVEMENT

Autophagy is an adaptive cellular process in PDL cells and osteocytes that facilitates managing mechanical stress. By promoting cell survival, regulating inflammation, and maintaining tissue stability, autophagy supports the biological adjustment required during orthodontic force application.[31,32] Extracellular matrix remodelling, particularly collagen turnover, enables periodontal tissues to adapt to orthodontic forces. Matrix receptors, metalloproteinases, and collagen internalisation pathways dismantle and reconstruct the matrix, consequently facilitating tooth movement and tissue repair.[33,34]

OTM is all about the precise interplay between immune and bone cells. Macrophages and T cells regulate osteoclastogenesis and direct bone remodelling via regulatory signals, highlighting the crucial role of the immune system.[35,36]

Circadian rhythm genes are involved in the regulation of bone metabolism and inflammation and so can have an effect on the rate and predictability of orthodontic outcome.[37] Hormones, metabolic diseases and medication can all have a significant effect on the tissue response to orthodontic force.[38]

TRANSLATIONAL AND CLINICAL IMPLICATIONS

Advances in the biological understanding of OTM have enabled translational strategies to accelerate tooth movement, including local delivery of cytokines or growth factors, minimally invasive surgical techniques, and optimisation of force systems.[39-41] Innovations in biomaterials and appliance design also aim to enhance biocompatibility and treatment efficacy. Upcoming research should integrate immunology and mechanobiology in orthodontics, develop personalised orthodontic protocols based on biological markers, translate findings into evidence-based clinical applications, and conduct high-quality human clinical trials to validate biologic modulation strategies.

CONCLUSION

Orthodontic tooth movement is a complicated process involving mechanotransduction, inflammation, cellular signalling, and tissue remodelling. Recent advances emphasize the central roles of osteocytes, immune regulation, and novel molecular pathways. Continued integration of basic science and clinical research will support the development of biologically driven therapies that are efficient, predictable, and patient-focused.

References

1. Burstone CJ. The biomechanics of tooth movement. *In*: Kraus BS, Riedel RA, editors. *Vistas in Orthodontics*. Philadelphia: Lea & Febiger; 1962.
2. Masella RS, Meister M. Current concepts in the biology of orthodontic tooth movement. *Am J Orthod Dentofacial Orthop*. 2006;129:458–468.
3. Davidovitch Z. Tooth movement. *Crit Rev Oral Biol Med*. 1991;2:411–450.
4. Krishnan V, Davidovitch Z. On a path to unfolding the biological mechanisms of orthodontic tooth movement. *J Dent Res*. 2009;88:597–608.
5. Li Y, Jacox LA, Little SH, Ko CC. Orthodontic tooth movement: the biology and clinical implications. *Kaohsiung J Med Sci*. 2018;34:207–214.
6. Vignery A. Osteocytes and bone remodeling. *Bone*. 2018;120:180–189.
7. Weltman B, Vig KW, Fields HW, Shanker S, Kaizar EE. Root resorption associated with orthodontic tooth movement. *Am J Orthod Dentofacial Orthop*. 2010;137:462–476.
8. Maltha JC, Kuijpers-Jagtman AM. Root resorption and orthodontic tooth movement. *Semin Orthod*. 2001;7:151–159.
9. Reitan K. Tissue rearrangement during retention of orthodontically rotated teeth. *Angle Orthod*. 1959;29:105–113.
10. Rygh P. Ultrastructural changes in pressure zones of human periodontium. *Angle Orthod*. 1973;43:271–285.
11. Kanzaki H, Chiba M, Shimizu Y, Mitani H. Periodontal ligament cells under mechanical stress induce osteoclastogenesis. *J Bone Miner Res*. 2002;17:210–220.
12. Schwarz AM. Tissue changes incident to orthodontic tooth movement. *Int J Orthod*. 1932;18:331–352.
13. Frost HM. Bone remodeling dynamics. Springfield: Charles C Thomas; 1963.
14. Baron R. Anatomy and ultrastructure of bone. *In*: Favus MJ, editor. *Primer on the Metabolic Bone Diseases*. 6th ed.
15. Klein-Nulend J, Bakker AD, Bacabac RG, Vatsa A, Weinbaum S. Mechanosensation and transduction in osteocytes. *Bone*. 2013;54:182–190.
16. Bonewald LF. The amazing osteocyte. *J Bone Miner Res*. 2011;26:229–238.

17. Robling AG, Turner CH. Mechanical signaling for bone modeling and remodeling. *Crit Rev Eukaryot Gene Expr*. 2009;19:319–338.
18. Nakashima T, Hayashi M, Fukunaga T, et al. Evidence for osteocyte regulation of bone homeostasis. *Nat Med*. 2011;17:1231–1234.
19. O'Brien CA. Control of RANKL gene expression. *Bone*. 2010;46:911–919.
20. Robling AG, Niziolek PJ, Baldridge LA, et al. Mechanical stimulation of bone in vivo reduces osteocyte expression of sclerostin. *J Biol Chem*. 2008;283:5866–5875.
21. Burger EH, Klein-Nulend J. Mechanotransduction in bone. *Bone*. 1999;25:127–130.
22. Theoleyre S, Wittrant Y, Tat SK, Fortun Y, Redini F, Heymann D. The molecular triad RANK/RANKL/OPG. *Endocr Rev*. 2004;25:207–227.
23. Kanzaki H, Chiba M, Takahashi I, Haruyama N, Nishimura M, Mitani H. Local RANKL gene transfer to periodontal tissue accelerates tooth movement. *Gene Ther*. 2006;13:678–685.
24. Udagawa N, Takahashi N, Yasuda H, et al. Osteoprotegerin produced by osteoblasts is an important regulator in osteoclast development. *Endocrinology*. 2000;141:3478–3484.
25. Yamaguchi M, Kasai K. Inflammation in periodontal tissues in response to orthodontic forces. *Orthod Craniofac Res*. 2005;8:234–242.
26. Alhashimi N, Frithiof L, Brudvik P, Bakhiet M. Orthodontic tooth movement and cytokine expression. *Am J Orthod Dentofacial Orthop*. 2001;119:301–309.
27. Grieve WG, Johnson GK, Moore RN, Reinhardt RA, DuBois LM. Prostaglandin E_2 and IL-1β levels in gingival crevicular fluid. *Am J Orthod Dentofacial Orthop*. 1994;105:369–374.
28. Baron R, Kneissel M. WNT signaling in bone homeostasis. *Bone*. 2013;56:246–253.
29. Poole KE, van Bezooijen RL, Loveridge N, et al. Sclerostin is a delayed secreted product of osteocytes. *Bone*. 2005;37:324–334.
30. Kook SH, Jang YS, Lee JC. Involvement of MAPK signaling in orthodontic tooth movement. *J Cell Biochem*. 2011;112:1510–1518.
31. Li J, Zhao Z, Yang J, et al. Autophagy in periodontal ligament cells under mechanical stress. *J Periodontal Res*. 2019;54:1–9.
32. Yang Y, Chen Y, Zhang C. Autophagy in orthodontic tooth movement. *Bone Res*. 2021;9:1–10.
33. Redlich M, Reichenberg E, Harari D, Zaks B, Shoshan S, Palmon A. The effect of mechanical force on collagen turnover. *J Dent Res*. 2001;80:1282–1286.
34. Holmbeck K, Bianco P, Pidoux I, et al. The role of matrix metalloproteinases in bone remodeling. *J Bone Miner Res*. 1999;14:476–486.
35. Takayanagi H. Osteoimmunology. *Nat Rev Immunol*. 2007;7:292–304.
36. Kiesel J, Kohl A, Lux CJ, Proff P. Immune response in orthodontic tooth movement. *Prog Orthod*. 2017;18:1–7.
37. Kawai M, Rosen CJ. The circadian clock and bone metabolism. *Clin Rev Bone Miner Metab*. 2010;8:31–42.
38. Krishnan V, Davidovitch Z. Biological mechanisms of tooth movement. *Semin Orthod*. 2015;21:3–12.
39. Alikhani M, Raptis M, Zoldan B, et al. Effect of micro-osteoperforations on orthodontic tooth movement. *Am J Orthod Dentofacial Orthop*. 2013;144:639–648.
40. Teixeira CC, Khoo E, Tran J, Chartres I, Liu Y, Thant LM. Cytokine expression and accelerated tooth movement. *J Dent Res*. 2010;89:1135–1141.
41. Wilcko WM, Wilcko MT, Bouquot JE, Ferguson DJ. Rapid orthodontics with alveolar reshaping. *Int J Periodontics Restorative Dent*. 2001;21:9–19.