POSTPARTUM DEPRESSION AND ITS IMPACT ON BREASTFEEDING, MOTHER-INFANT BONDING AND CHILD DEVELOPMENT

Authored and Submitted by

FIROJ KHAN

Research Scholar, Department of Health Science ARNI UNIVERSITY

Indora, Kathgarh, District Kangra, Himachal Pradesh, India

Under the Supervision of

(Dr.) CharanPreet Singh , Professor

Research Supervisor, Department of Health Science

Received:Oct 01, 2026 Accepted:Oct 29, 2026 Published:Nov 01, 2026

Abstract:

Postpartum depression (PPD) is a clinically important depressive condition arising during pregnancy or after childbirth and extending beyond the brief, self-limiting emotional changes commonly called the baby blues. It affects maternal well-being while also influencing feeding decisions, responsive caregiving and the relational environment in which early development occurs. This paper synthesises contemporary evidence on the connections between postpartum depressive symptoms, breastfeeding, mother-infant bonding and child development, with particular attention to the Indian public-health context. The evidence indicates a bidirectional relationship: depressive symptoms can undermine breastfeeding confidence, initiation and continuation, while painful or unsuccessful breastfeeding experiences can intensify distress. PPD is also associated with impaired bonding, reduced maternal sensitivity and less consistent contingent interaction. Through these pathways, persistent or severe symptoms may contribute to

difficulties in infant sleep, growth, emotional regulation, cognition and language; however, outcomes are probabilistic rather than inevitable and are shaped by social support, poverty, violence, obstetric complications, infant illness and access to care. The paper proposes an integrated prospective cohort design using the Edinburgh Postnatal Depression Scale, validated bonding and breastfeeding measures, and age-appropriate developmental screening. It recommends repeated screening linked to referral, breastfeeding assistance that protects maternal choice, family-inclusive psychosocial support, stepped care and follow-up through antenatal, postnatal, immunisation and child-health platforms. The central argument is that maternal mental health, breastfeeding support and early childhood development should be addressed as one interconnected continuum rather than as separate programmes.

Keywords: postpartum depression; breastfeeding; maternal-infant bonding; child development; maternal mental health; India; public health.

1. Introduction

1.1 Background:

Childbirth is often represented as an exclusively joyful event, yet the transition to motherhood can involve physical pain, sleep disruption, changing relationships, financial pressure and an abrupt increase in responsibility. Emotional distress in this period does not indicate weak character or poor motherhood. It can reflect the interaction of biological vulnerability, previous mental illness, obstetric experiences and social conditions. When depressive symptoms persist, intensify or interfere with daily functioning, they require respectful clinical attention.

The World Health Organization estimates that mental disorders affect approximately 13% of women after childbirth worldwide, with a higher burden in developing-country settings. Indian studies report considerable variation because populations, timing and screening thresholds differ. Recent reviews nevertheless show that PPD is common and frequently overlooked. A synthesis of Indian evidence has reported estimates ranging widely across settings, while a 2024 review cited a pooled prevalence near 22%. These figures should be read as screening-based estimates rather than a single fixed national rate (World Health Organization, n.d.; Panolan et al., 2024; Rajeev et al., 2024).

PPD matters beyond symptom counts. A mother may struggle to initiate or maintain breastfeeding, experience guilt when feeding does not proceed as expected, withdraw from interaction, or find it difficult to interpret and respond to infant cues. These experiences can affect bonding and the daily exchanges through which infants develop security, communication and self-regulation. At the same time, supportive partners, responsive health workers and timely treatment can protect both mother and child. The relationship is therefore dynamic, modifiable and embedded in family and health-system contexts.

1.2 Problem statement:

Maternal health services in India have expanded institutional delivery and routine contacts, but emotional well-being is not always assessed systematically. Women may hide sadness, intrusive thoughts or feelings of inadequacy because of stigma, fear of judgement, limited privacy or the belief that suffering is a normal part of motherhood. Breastfeeding difficulties may be addressed only as technical feeding problems, without recognising depression, anxiety, pain or coercive family expectations. Similarly, child-development concerns may be considered separately from maternal functioning and the quality of early interaction. Fragmented care can therefore miss a linked pathway of risk.

1.3 Aim and objectives:

Aim: To examine postpartum depression and its relationship with breastfeeding practices, mother-infant bonding and early child development, and to propose an integrated public-health response.

- To describe the concept, burden, clinical presentation and determinants of postpartum depression.
- To examine the association between postpartum depressive symptoms and breastfeeding initiation, exclusivity, continuation and maternal feeding experience.
- To assess how postpartum depression may influence mother-infant bonding and responsive caregiving.
- To synthesise evidence on developmental outcomes associated with maternal postpartum depression.
- To identify protective factors, moderators and equity concerns affecting these relationships.

- To propose a feasible research methodology and evidence-informed recommendations for maternal and child health services.

1.4 Research questions:

- What maternal, infant, family and health-system factors are associated with postpartum depressive symptoms?
- How are the severity and duration of depressive symptoms related to breastfeeding practices and feeding confidence?
- Does impaired bonding mediate or explain part of the relationship between PPD and child-development outcomes?
- Which forms of family, peer and professional support protect breastfeeding, bonding and development?
- Do relationships differ by rural residence, socioeconomic position, parity, infant sex, disability or access to care?

2. Conceptual and Clinical Foundations:

2.1 Definition and distinction from baby blues and psychosis:

Postpartum depression involves a sustained pattern of depressed mood or loss of interest accompanied by symptoms such as guilt, hopelessness, altered sleep or appetite beyond infant-related disruption, impaired concentration, psychomotor change, fatigue and thoughts of death or self-harm. Onset may occur during pregnancy or following birth. Baby blues are milder and usually resolve within about two weeks. Postpartum psychosis is rare but constitutes an emergency, especially when there is severe confusion, mania, delusion, hallucination or risk to the mother or infant. Screening instruments identify probable cases; they do not replace diagnostic assessment.

2.2 Determinants and risk architecture:

Level	Illustrative determinants	Public-health implication
Maternal	Previous depression or anxiety; trauma; unplanned pregnancy; anaemia; chronic illness; sleep deprivation	Screen during pregnancy and repeatedly after birth; review medical and psychosocial needs.
Obstetric/infant	Complicated birth; caesarean delivery; pain; preterm birth; neonatal illness; feeding difficulty	Link maternity, newborn, lactation and mental-health services.
Family	Low partner support; conflict; intimate-partner violence; son preference; unequal care burden	Use confidential enquiry, safety pathways and family education without compromising autonomy.
Socioeconomic	Poverty; food insecurity; insecure employment; migration; transport barriers	Combine clinical care with social protection and community follow-up.
Health system	Short visits; stigma; lack of trained staff; screening without referral; fragmented records	Adopt stepped care, referral tracking and supervision.

2.3 Conceptual pathway:

The proposed framework treats PPD as both an outcome of accumulated vulnerability and an exposure affecting caregiving. Structural disadvantage and stressful events influence depressive symptoms. Symptoms may then affect energy, confidence, decision-making and sensitivity to infant cues. These changes can disrupt breastfeeding and bonding, which may influence nutrition, stress regulation and opportunities for learning. The pathway is not linear: infant illness or breastfeeding pain can worsen maternal mood, and supportive care can interrupt the cycle at several points.

3. Review of Evidence:

3.1 Postpartum depression and breastfeeding:

The association between PPD and breastfeeding is best understood as bidirectional. Depressive symptoms may reduce confidence, make night-time feeding more difficult, complicate

interpretation of infant hunger cues and reduce persistence when pain or low milk perception occurs. Women may also receive contradictory advice and experience intense pressure to breastfeed, turning feeding into a source of guilt. Conversely, early cessation, unresolved pain, perceived insufficient milk and a mismatch between intended and actual feeding can contribute to distress. Accordingly, a simple statement that breastfeeding prevents depression, or that depression causes breastfeeding failure, overstates the evidence.

Observational research commonly associates PPD with delayed initiation, reduced exclusivity and shorter duration, although effect sizes vary and residual confounding is likely. A 2024 Nigerian study reported associations between PPD, poor maternal-infant bonding and late breastfeeding initiation. Other contemporary evidence indicates that breastfeeding, maternal mood and interaction influence one another. Clinical care should therefore address mood, pain, latch, infant growth, medication questions, sleep and maternal preference together rather than measuring exclusivity as the only success criterion (Sanni et al., 2024; Coo et al., 2024; Carvalho Hilje et al., 2024).

3.2 Postpartum depression and mother-infant bonding:

Bonding refers to the mother's emotional orientation towards her infant, whereas attachment describes the child's developing relationship with a caregiver. Depression may be accompanied by emotional numbness, irritability, self-criticism or fear of causing harm. These experiences can make closeness feel effortful and may reduce pleasure in interaction. Some mothers remain highly responsive despite severe internal distress, so bonding difficulty should never be assumed from diagnosis alone.

Across studies, higher depressive symptoms are generally associated with more bonding difficulties and lower observed sensitivity. The relationship can also operate in reverse: persistent feelings of disconnection may reinforce guilt and depressive thinking. Recent network and observational studies support this reciprocal interpretation. Assessment should ask about enjoyment, closeness, avoidance, anger, fear and caregiving confidence in a non-judgemental manner (Qi et al., 2025; Harasawa et al., 2025).

3.3 Postpartum depression and child development:

Infant and child outcomes associated with persistent maternal depression include differences in emotional regulation, sleep, behaviour, language and cognitive performance. Several mechanisms may contribute: reduced contingent interaction, fewer language-rich exchanges, disrupted routines, feeding difficulties, household stress and biological stress pathways. Severe poverty, violence, prematurity and paternal depression may independently affect the same outcomes and must be considered in analysis.

Association does not mean that every child of a depressed mother will experience delay. Many children develop well because symptoms are brief, treatment is effective or other caregivers provide stable and responsive care. The most concerning pattern is sustained, recurrent or severe depression combined with limited support and multiple adversities. Reviews have consistently emphasised effects across maternal health, partner relationships and child development, while newer cohort evidence suggests particular vulnerability in socio-emotional, language and fine-motor domains (Slomian et al., 2019; Alfayumi-Zeadna et al., 2025).

3.4 Protective factors and moderators:

- Early recognition and access to acceptable psychological or psychiatric care.
- Practical and emotional partner support, including shared night-time and domestic responsibilities.
- Skilled breastfeeding support that respects informed maternal choice and avoids blame.
- Continuity with a trusted health worker and confidential communication.
- Adequate nutrition, sleep opportunities, pain control and management of maternal illness.
- Responsive involvement of another caregiver when the mother is unwell.
- Social protection, safe housing and intervention for intimate-partner violence.

3.5 Evidence limitations:

Much of the literature is observational and relies on self-reported screening measures. Studies use different postpartum windows, thresholds and definitions of breastfeeding and bonding. Reverse causation is common, and unmeasured confounding can produce apparently direct effects. Cross-sectional designs cannot establish temporal ordering. Cultural adaptation of scales, inclusion of

women with low literacy, and representation of adolescents, migrants, women with disabilities and mothers of sick infants remain inconsistent. These limitations justify prospective, repeated-measure designs and careful causal language.

4. Proposed Methodology:

4.1 Study design and setting:

A prospective mixed-method cohort study is proposed. Women would be enrolled during the third trimester or within seven days after childbirth and followed at approximately 6 weeks, 3 months, 6 months and 12 months postpartum. Recruitment may occur through selected public and private facilities and their community catchment areas in Kangra district and, if approved, additional sites that improve rural and socioeconomic representation. The design permits examination of change, temporal ordering and service pathways while qualitative interviews explain experiences hidden by scale scores.

4.2 Population, eligibility and sample:

Eligible participants would be mothers aged 18 years or older with a live birth who reside in the study area and intend to remain for twelve months. Mothers of preterm or medically vulnerable infants should not be automatically excluded; instead, infant condition should be measured because it is clinically important. Women unable to consent at recruitment or requiring emergency medical or psychiatric care should receive immediate care and may be approached later only when ethically appropriate. A sample of approximately 500 mother-infant dyads would permit multivariable analysis and allowance for attrition, but the final sample must follow a documented power calculation based on the primary outcome and expected prevalence.

4.3 Measures

Domain	Proposed measure	Timing
Depressive symptoms	Edinburgh Postnatal Depression Scale (EPDS), culturally validated language version; diagnostic confirmation for a subsample where feasible	Baseline/6 weeks/3/6/12 months
Breastfeeding	Early initiation, exclusive breastfeeding, current feeding, duration, pain, difficulty, intention-experience mismatch and Breastfeeding Self-Efficacy Scale-Short Form	Birth record and every follow-up
Bonding	Postpartum Bonding Questionnaire or Mother-to-Infant Bonding Scale; optional structured interaction observation	6 weeks/3/6 months
Child development	Age-appropriate validated developmental screening covering communication, gross and fine motor, problem-solving and socio-emotional domains	6 and 12 months
Covariates	Age, parity, education, wealth, caste/social group, residence, pregnancy intention, obstetric morbidity, infant illness, violence, partner support, sleep and prior mental illness	Baseline and updated follow-up

Qualitative experience	Semi-structured interviews with purposively selected mothers, partners and health workers	3-9 months

4.4 Data collection and quality assurance:

Female investigators trained in respectful maternity and mental-health interviewing should collect data privately. Translation should use forward translation, expert reconciliation, back-translation and cognitive interviewing. Electronic forms should include range checks without forcing inaccurate responses. Ten per cent of non-sensitive records may be verified through supervised checks. Referral information must be available before screening begins. Instruments should be piloted outside the final sample, and internal consistency, missingness and floor or ceiling effects should be reported rather than relying on a single reliability coefficient.

4.5 Statistical analysis:

Analysis should report prevalence with 95% confidence intervals at each wave, trajectories of symptom severity and attrition patterns. Breastfeeding outcomes may be analysed with modified Poisson or logistic regression, while duration can be examined using survival analysis. Continuous bonding and development scores may be modelled using linear mixed-effects models; clinically defined developmental risk may use generalized mixed models. Confounders must be selected from a directed conceptual model rather than stepwise significance testing. Mediation analysis may examine whether bonding or feeding experience explains part of an association between PPD and later development, but only when exposure precedes mediator and outcome. Sensitivity analyses should address prior antenatal depression, infant illness, missing data and alternative EPDS thresholds. Results should present adjusted estimates, confidence intervals and absolute risks.

4.6 Qualitative analysis and integration:

Interviews should explore emotional change, feeding expectations, family responses, stigma, help-seeking, interactions with health workers and the meaning of bonding. Reflexive thematic analysis may be used with an audit trail, diverse coding perspectives and attention to negative cases. Integration should occur through joint displays that compare statistical patterns with lived

explanations—for example, why two mothers with similar depression scores have different breastfeeding experiences because one receives practical support and the other faces criticism or violence.

4.7 Ethical considerations:

Ethics approval and administrative permission must be obtained before recruitment. Consent should be voluntary, understandable and separate from clinical care. Privacy is essential because disclosure of depression, self-harm thoughts or violence may create risk. A predefined safety protocol should cover immediate risk assessment, supportive response, referral, emergency escalation and documented follow-up. Screening without a functioning referral pathway is ethically inadequate. Data should be de-identified, encrypted and retained only for the approved period. Infant developmental screening should be explained as an initial assessment, not a diagnosis, with referral for concerning results.

5. Expected Findings and Interpretation:

The study is expected to identify a subgroup with persistent or recurrent depressive symptoms and to demonstrate that risk is socially patterned. Greater symptom severity is likely to be associated with lower breastfeeding self-efficacy, more reported feeding difficulties and shorter breastfeeding duration. Depressive symptoms are also expected to correlate with bonding difficulty, although substantial variation will remain. Developmental risk may be greater when maternal symptoms are persistent and co-occur with poverty, infant illness, violence or low support.

These expectations are hypotheses, not fabricated results. The final paper must replace this section with analysed findings, including participant flow, missing-data assessment, descriptive tables, model diagnostics and confidence intervals. A statistically significant association should not be interpreted automatically as clinically important, and a non-significant finding should not be equated with absence of effect when precision is low.

Hypothesised pathway	Expected direction	Interpretive caution
PPD severity -> breastfeeding confidence	Negative	Mood and feeding difficulty may influence each other.
PPD persistence -> bonding quality	Negative	Internal distress does not always produce observable disengagement.
Bonding difficulty -> later developmental risk	Positive	Infant illness and social adversity may confound the pathway.
Family/professional support x PPD	Protective moderation	Support quality, control and timing must be measured.
PPD -> development through feeding/bonding	Partial mediation	Requires longitudinal ordering; causal claims remain cautious.

6. Discussion:

6.1 Integrated interpretation:

The reviewed evidence supports an integrated view of postpartum depression as a maternal disorder with relational and developmental implications. Breastfeeding, bonding and child development should not be placed in a moral hierarchy. A mother receiving treatment or using mixed or replacement feeding can still form a warm relationship and support healthy development. Public-health practice should focus on safe nutrition, responsive care, maternal recovery and informed choice rather than treating one feeding pattern as proof of good motherhood.

The strongest interpretation is not that PPD directly determines a child's future, but that it can alter a chain of everyday opportunities: recognising hunger, sustaining eye contact, talking and playing, attending follow-up, and seeking help when the infant is unwell. Each link is influenced by family resources and health-system responsiveness. This creates multiple intervention points and avoids blaming the mother for structural conditions.

6.2 Implications for India and Himachal Pradesh:

In geographically dispersed settings, travel time, privacy in multigenerational households and scarcity of specialists may limit access. Maternal mental-health support can be embedded within antenatal care, delivery discharge, home visits, immunisation sessions and child-development contacts. ASHAs, ANMs, nurses and medical officers can contribute to identification and follow-up when they receive training, supervision, manageable workloads and clear referral options. Digital follow-up may help some women but should not replace confidential in-person care or assume personal phone access.

6.3 Clinical and programme recommendations:

- Introduce repeated, validated screening during pregnancy and postpartum, with immediate review of self-harm responses.
- Link every screening programme to stepped referral, treatment availability and follow-up documentation.
- Train maternity and child-health teams to discuss mood, sleep, pain, feeding and bonding without blame.
- Provide early lactation assistance while protecting the mother's informed feeding choice and medication needs.
- Include partners or trusted relatives with the mother's consent, focusing on practical workload-sharing and emotional support.
- Integrate developmental surveillance and responsive-care counselling into routine infant contacts.
- Establish pathways for intimate-partner violence, severe mental illness, postpartum psychosis and safeguarding emergencies.

- Audit coverage and outcomes by rurality, poverty, disability, language, caste/social group and facility type.

7. Conclusion:

Postpartum depression is common, treatable and consequential. Its relationship with breastfeeding is reciprocal, its association with bonding is meaningful but not deterministic, and its influence on child development is shaped by symptom persistence and the wider caregiving environment. The most humane and scientifically defensible response is integrated care that supports maternal recovery, safe infant feeding, sensitive interaction and developmental follow-up. Research should use longitudinal measurement, culturally validated instruments, explicit confounder control and cautious causal interpretation. Services should make help available before asking women to disclose distress. When health systems listen without judgement and families share the work of care, the pathway from depression to adverse maternal and child outcomes can be interrupted.

References:

American College of Obstetricians and Gynecologists. (2023a). Screening and diagnosis of mental health conditions during pregnancy and postpartum. Obstetrics & Gynecology, 141(6), 1232-1261. https://doi.org/10.1097/AOG.0000000000005200

American College of Obstetricians and Gynecologists. (2023b). Treatment and management of mental health conditions during pregnancy and postpartum. Obstetrics & Gynecology, 141(6), 1262-1288.

Alfayumi-Zeadna, S., et al. (2025). The impact of postpartum depression on infant developmental outcomes. Journal of Affective Disorders.

Brockington, I. F., Oates, J., George, S., Turner, D., Vostanis, P., Sullivan, M., Loh, C., & Murdoch, C. (2001). A screening questionnaire for mother-infant bonding disorders. Archives of Women's Mental Health, 3, 133-140.

Carvalho Hilje, C., et al. (2024). The role of breastfeeding and formula feeding regarding postpartum depression and mother-child bonding. Scientific Reports, 14.

Coo, S., et al. (2024). Quality of mother-infant interaction, breastfeeding, and maternal mental health. Infant Behavior and Development, 75, 101935.

Cox, J. L., Holden, J. M., & Sagovsky, R. (1987). Detection of postnatal depression: Development of the 10-item Edinburgh Postnatal Depression Scale. British Journal of Psychiatry, 150(6), 782-786.

Dennis, C.-L. (2003). The Breastfeeding Self-Efficacy Scale: Psychometric assessment of the short form. Journal of Obstetric, Gynecologic & Neonatal Nursing, 32(6), 734-744.

Harasawa, N., et al. (2025). A network analysis of postpartum depression and mother-infant bonding. Communications Psychology, 3.

Howard, L. M., & Khalifeh, H. (2020). Perinatal mental health: A review of progress and challenges. World Psychiatry, 19(3), 313-327.

Kroenke, K., Spitzer, R. L., & Williams, J. B. W. (2001). The PHQ-9: Validity of a brief depression severity measure. Journal of General Internal Medicine, 16, 606-613.

Norhayati, M. N., Hazlina, N. H. N., Asrenee, A. R., & Emilin, W. M. A. W. (2015). Magnitude and risk factors for postpartum symptoms: A literature review. Journal of Affective Disorders, 175, 34-52.

Panolan, S., et al. (2024). Prevalence and associated risk factors of postpartum depression in India: A comprehensive review. Annals of Indian Psychiatry, 8(1).

Qi, W., et al. (2025). Postpartum depression and maternal-infant bonding: A systematic investigation. BMC Pregnancy and Childbirth, 25.

Rajeev, S. P., et al. (2024). India's silent struggle: A scoping review on postpartum depression. Cureus, 16.

Sanni, S. O., et al. (2024). Influence of postpartum depression on maternal-infant bonding and breastfeeding initiation. BMC Pregnancy and Childbirth, 24.

Slomian, J., Honvo, G., Emonts, P., Reginster, J.-Y., & Bruyère, O. (2019). Consequences of maternal postpartum depression: A systematic review of maternal and infant outcomes. Women's Health, 15, 1-55.

Stewart, D. E., & Vigod, S. N. (2019). Postpartum depression: Pathophysiology, treatment, and emerging therapeutics. Annual Review of Medicine, 70, 183-196.

UNICEF. (2021). Caring for the caregiver: Implementing the nurturing care framework. United Nations Children's Fund.

World Health Organization. (2016). mhGAP intervention guide for mental, neurological and substance use disorders in non-specialized health settings (Version 2.0).

World Health Organization. (2022). WHO guide for integration of perinatal mental health in maternal and child health services.

World Health Organization. (2023). Guideline for complementary feeding of infants and young children 6-23 months of age.

World Health Organization. (n.d.). Perinatal mental health. https://www.who.int/teams/mental-health-and-substance-use/promotion-prevention/perinatal-mental-health

World Health Organization, UNICEF, & World Bank Group. (2018). Nurturing care for early childhood development: A framework for helping children survive and thrive to transform health and human potential.

About the Author:

Authored and Submitted by

FIROJ KHAN

Research Scholar, Department of Health Science ARNI UNIVERSITY

Indora, Kathgarh, District Kangra, Himachal Pradesh, India

Under the Supervision of

(Dr.) CharanPreet Singh , Professor

Research Supervisor, Department of Health Science